

NAS5-96101  
INTERIM  
70-75-112  
O.T.  
0-7238

Interim Contractor Report #2

12/01/96 - 12/31/96

**Physics of Boundaries and their Interactions in Space Plasmas**

Dr. Nojan Omid  
Dr. Homayoun Karimabadi  
Dr. Dietmar Krauss-Varban

SciberNet, Inc.  
5414 Oberlin Drive, Suite 251  
San Diego, CA 92121

## **I. Bow Shock & Foreshock:**

We are continuing our large scale hybrid simulations of the bow shock. In particular we have examined the effect of IMF direction on the large scale structure of the shock/Foreshock system. By performing shock normal diagnostics, we were able to demonstrate that the foreshock is the result of ion reflection and leakage from only the quasi-parallel side of the bow shock with shock normal angles below about 42 degrees. We also spent a considerable time preparing diagnostics and figures in order to develop a coherent picture of our simulations results which were then presented at the Fall American Geophysical Union meeting.

## **II. Kelvin-Helmholtz Instability at the Magnetopause and the Magnetotail:**

We used a combination of 2-D and 3-D hybrid simulations to examine the effect of the velocity shear on the stability of the magnetotail and the magnetopause. We demonstrated that the presence of secondary ions at the magnetopause (e.g., cold ring ions, heavy elements) can lead to the excitation of Kelvin Helmholtz (KH) instability at or near the subsolar point. This is in sharp contrast to previous studies where it was thought the KH could only be excited at the flanks. We also identified several ways that KH could be excited in the magnetotail. Our preliminary results suggest that the KH could lead to destabilization of the tearing mode in the magnetotail. The results were presented at the Fall American Geophysical Union meeting.

## **III. Large-Scale Hybrid Simulations of the Magnetotail:**

Previously, we successfully used the 2-D hybrid code to simulate the large scale structure of the distant magnetotail. As a next step in our research, we have started tackling the more complex physical configuration of near-Earth reconnection and its consequences. The near-Earth problem has several added complications: (1) such simulations require a 2-D equilibrium model; (2) a reflecting particle boundary condition is needed at the earthward side; (3) open (self-adjusting) boundary conditions should be used at the lobes to allow the internal plasma to determine the reconnection time evolution. We have implemented these changes to our code and have for the first time performed several large scale simulations with these new boundary conditions, the results of which were presented at the Fall American Geophysical Union meeting.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 01/07/97	3. REPORT TYPE AND DATES COVERED Interim Contractor Report, 12/01/96-12/31/96		
4. TITLE AND SUBTITLE Physics of Boundaries and their Interactions in Space Plasmas		5. FUNDING NUMBERS C: NAS5-96101		
6. AUTHOR(S) Dr. Nojan Omidi, Dr. Homayoun Karimabadi, and Dr. Dietmar Krauss-Varban				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) SciberNet, Inc. 5414 Oberlin Drive, Suite 251 San Diego, CA 92121		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Goddard Space Flight Center Greenbelt, MD 20771		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Unclassified-Unlimited		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words)  <p>This report describes the work done by SciberNet, Inc. during the month of December. Much of the work performed during this month involved preparing the three talks that we gave at the Fall American Geophysical Union meeting. To this end, we further developed and refined our demonstration of the fact that the quasi-parallel part of the bow shock is the sole contributor to the ion beams in the foreshock. In a second area of research, we identified new mechanisms for the excitation of the Kelvin-Helmholtz instability at the magnetopause and the magnetotail. Given the 3-D nature of our simulations, we have also started development of data visualization tools in 3-D. Finally, we made several large scale 2-D hybrid simulations of the near magnetotail with new boundary conditions to allow for unbiased plasma and flow evolution during magnetic reconnection.</p>				
14. SUBJECT TERMS Foreshock, bow shock, magnetotail, reconnection, kinetic simulations		15. NUMBER OF PAGES 1		
		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	